

[illegible]

4 wherein

5 A is an electrophilic group that reacts with a cysteine residue of said RdRp  
6 protein;

7 Ar<sup>1</sup> is a member selected from the group consisting of substituted or  
8 unsubstituted aryl and substituted or unsubstituted heteroaryl;

9 X<sup>1</sup> is a member selected from the group consisting of -H, substituted or  
10 unsubstituted alkyl, substituted or unsubstituted aryl, substituted or unsubstituted  
11 heteroaryl, -CN, -CO<sub>2</sub>H, -SO<sub>3</sub>H, -C(O)NHOH, -NH<sub>2</sub>, -OH, -NH(lower alkyl), -O(lower  
12 alkyl), -N(lower alkyl)<sub>2</sub>, and -C(O)-NH(3-tetrazolyl);

13 L<sup>1</sup> is a divalent linking group selected from the group consisting of  
14 -CH<sub>2</sub>CH<sub>2</sub>-, -CH=CH-, -C≡C-, -O-, -S(O)<sub>n</sub>-, -N(R<sub>a</sub>)-, -C(O)-, -C(O)O-, -SO<sub>2</sub>N(R<sub>a</sub>)-,  
15 -CON(R<sub>a</sub>)-, -N(R<sub>a</sub>)CON(R<sub>b</sub>)-, -N(R<sub>a</sub>)N(R<sub>b</sub>)-, -N(R<sub>a</sub>)SO<sub>2</sub>N(R<sub>b</sub>)-, -N(R<sub>a</sub>)SO<sub>2</sub>-, -N(R<sub>a</sub>)-O-,  
16 =N-O-, lower alkylene, -O-lower alkylene, -S(O)<sub>n</sub>-lower alkylene, N(R<sub>a</sub>)-lower alkylene,  
17 -SO<sub>2</sub>N(R<sub>a</sub>)-lower alkylene, lower alkylene-SO<sub>2</sub>N(R<sub>a</sub>)-, -CON(R<sub>a</sub>)-lower alkylene, lower  
18 alkylene-CON(R<sub>a</sub>)-, -N(R<sub>a</sub>)CON(R<sub>b</sub>)-lower alkylene, lower alkylene-N(R<sub>a</sub>)N(R<sub>b</sub>)-,  
19 -N(R<sub>a</sub>)SO<sub>2</sub>N(R<sub>b</sub>)-lower alkylene, -N(R<sub>a</sub>)-O-lower alkylene, lower alkylene-N(R<sub>a</sub>)-O-,  
20 =N-O-lower alkylene, lower heteroalkylene, -O-lower heteroalkylene, -S(O)<sub>n</sub>-lower  
21 heteroalkylene, N(R<sub>a</sub>)-lower heteroalkylene, -SO<sub>2</sub>N(R<sub>a</sub>)-lower heteroalkylene, lower  
22 heteroalkylene-SO<sub>2</sub>N(R<sub>a</sub>)-, -CON(R<sub>a</sub>)-lower heteroalkylene, lower  
23 heteroalkylene-CON(R<sub>a</sub>)-, -N(R<sub>a</sub>)CON(R<sub>b</sub>)-lower heteroalkylene, lower  
24 heteroalkylene-N(R<sub>a</sub>)N(R<sub>b</sub>)-, -N(R<sub>a</sub>)SO<sub>2</sub>N(R<sub>b</sub>)-lower heteroalkylene, -N(R<sub>a</sub>)-O-lower  
25 heteroalkylene, lower heteroalkylene-N(R<sub>a</sub>)-O-, =N-O-lower alkylene, aryl and  
26 heteroaryl;

27 X<sup>2</sup> is a member selected from the group consisting of substituted or  
28 unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted  
29 cycloalkyl, and substituted or unsubstituted heterocycloalkyl;

30 L<sup>2</sup> is a divalent linking group selected from the group consisting of  
31 -CH<sub>2</sub>CH<sub>2</sub>-, -(C(R<sub>c</sub>)=C(R<sub>d</sub>))<sub>m</sub>-, -O-, -S(O)<sub>n</sub>-, -N(R<sub>e</sub>)-, -C(O)-, -C(O)O-, -SO<sub>2</sub>N(R<sub>e</sub>)-,  
32 -CON(R<sub>e</sub>)-, -N(R<sub>e</sub>)CON(R<sub>f</sub>)-, -N(R<sub>e</sub>)N(R<sub>f</sub>)-, -N(R<sub>e</sub>)SO<sub>2</sub>N(R<sub>f</sub>)-, -N(R<sub>e</sub>)-O-, =N-O-, lower  
33 alkylene, perfluoro lower alkylene, polyfluoro lower alkylene, -O-lower alkylene,  
34 -S(O)<sub>n</sub>-lower alkylene, N(R<sub>e</sub>)-lower alkylene, -SO<sub>2</sub>N(R<sub>e</sub>)-lower alkylene, lower  
35 alkylene-SO<sub>2</sub>N(R<sub>e</sub>)-, -CON(R<sub>e</sub>)-lower alkylene, lower alkylene-CON(R<sub>e</sub>)-,  
36 -N(R<sub>e</sub>)CON(R<sub>f</sub>)-lower alkylene, lower alkylene-N(R<sub>e</sub>)N(R<sub>f</sub>)-, -N(R<sub>e</sub>)SO<sub>2</sub>N(R<sub>f</sub>)-lower  
37 alkylene, -N(R<sub>e</sub>)-O-lower alkylene, lower alkylene-N(R<sub>e</sub>)-O-, =N-O-lower alkylene,

38 lower heteroalkylene, -O-lower heteroalkylene, -S(O)<sub>n</sub>-lower heteroalkylene, N(R<sub>e</sub>)-lower  
 39 heteroalkylene, -SO<sub>2</sub>N(R<sub>e</sub>)-lower heteroalkylene, lower heteroalkylene-SO<sub>2</sub>N(R<sub>e</sub>)-,  
 40 -CON(R<sub>e</sub>)-lower heteroalkylene, lower heteroalkylene-CON(R<sub>e</sub>)-, -N(R<sub>e</sub>)CON(R<sub>f</sub>)-lower  
 41 heteroalkylene, lower heteroalkylene-N(R<sub>e</sub>)N(R<sub>f</sub>)-, -N(R<sub>e</sub>)SO<sub>2</sub>N(R<sub>f</sub>)-lower  
 42 heteroalkylene, -N(R<sub>e</sub>)-O-lower heteroalkylene, lower heteroalkylene-N(R<sub>e</sub>)-O-,  
 43 =N-O-lower alkylene, aryl and heteroaryl, wherein R<sub>a</sub>, R<sub>b</sub>, R<sub>c</sub>, R<sub>d</sub>, R<sub>e</sub> and R<sub>f</sub> are each  
 44 members independently selected from the group consisting of H, lower alkyl, lower  
 45 heteroalkyl, -C(O)-lower alkyl, -C(O)-lower heteroalkyl, -S(O)<sub>2</sub>-lower alkyl, and -  
 46 S(O)<sub>2</sub>-lower heteroalkyl;

47 the subscript n is an integer of from 0 to 2;

48 the subscript m is an integer of from 0 to 3;

49 the bond between X<sup>2</sup> and A can be a single, double or triple bond,

50 depending on the nature of X<sup>2</sup> and A; and

51 wherein when L<sup>1</sup> and L<sup>2</sup> may be linked together *via* a single bond, -O-, -S-  
 52 or amide group to form a new 5 to 7 membered ring;

53 with the proviso that when A is an sp<sup>2</sup>-hybridized carbon atom and X<sup>2</sup> is substituted or  
 54 unsubstituted rhodanine, L<sup>1</sup> is not -CH<sub>2</sub>-CH<sub>2</sub>-, -CH=CH-, -C≡C- or aryl.

1 10. A compound in accordance with claim 9, wherein

2 X<sup>2</sup> is selected from the group consisting of a 5 to 7 membered cycloalkyl  
 3 ring, a 5 to 7 membered heterocycloalkyl ring containing from 1 to 3 heteroatoms, an aryl  
 4 group and a heteroaryl group;


5 A is selected from the group consisting of an sp<sup>2</sup>-hybridized carbon atom  
 6 and an sp<sup>3</sup>-hybridized carbon atom;

7 L<sup>2</sup> is a single bond; and

8 X<sup>2</sup> and A are joined *via* a single or double bond.

1 11. A compound in accordance with claim 10, wherein

2 X<sup>2</sup>-A- is selected from the group consisting of:




wherein  $R_g$  is selected from the group consisting of H, lower alkyl, lower alkoxy and F;  $R_h$  is selected from the group consisting of H,  $-S(O)_n$ -lower alkyl,  $-S(O)_n$ -lower heteroalkyl,  $-S(O)_n$ -aryl and  $-S(O)_n$ -heteroaryl;

W is CH or N; Hal is a halogen atom; and

$X^2$  is a substituted or unsubstituted member selected from the group consisting of a 5-6 membered cycloalkyl, 5-6 membered heterocycloalkyl containing from 1 to 3 heteroatoms, heteroaryl containing from 1 to 3 heteroatoms and aryl.

12. A compound having the formula (II):



wherein

$Ar^1$  is a member selected from the group consisting of substituted or unsubstituted aryl and substituted or unsubstituted heteroaryl;

$X^1$  is a member selected from the group consisting of -H, substituted or unsubstituted alkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, -CN,  $-CO_2H$ ,  $-SO_3H$ ,  $-C(O)NHOH$ ,  $-NH_2$ ,  $-OH$ ,  $-NH$ (lower alkyl),  $-O$ (lower alkyl),  $-N$ (lower alkyl)<sub>2</sub>, and  $-C(O)-NH$ (3-tetrazolyl);

$L^1$  is a divalent linking group selected from the group consisting of  $-CH_2CH_2-$ ,  $-CH=CH-$ ,  $-C\equiv C-$ ,  $-O-$ ,  $-S(O)_n-$ ,  $-N(R_a)-$ ,  $-C(O)-$ ,  $-C(O)O-$ ,  $-SO_2N(R_a)-$ ,  $-CON(R_a)-$ ,  $-N(R_a)CON(R_b)-$ ,  $-N(R_a)N(R_b)-$ ,  $-N(R_a)SO_2N(R_b)-$ ,  $-N(R_a)SO_2-$ ,  $-N(R_a)-O-$ ,  $=N-O-$ , lower alkylene,  $-O$ -lower alkylene,  $-S(O)_n$ -lower alkylene,  $N(R_a)$ -lower alkylene,  $-SO_2N(R_a)$ -lower alkylene, lower alkylene- $SO_2N(R_a)-$ ,  $-CON(R_a)$ -lower alkylene, lower alkylene- $CON(R_a)-$ ,  $-N(R_a)CON(R_b)$ -lower alkylene, lower alkylene- $N(R_a)N(R_b)-$ ,  $-N(R_a)SO_2N(R_b)$ -lower alkylene,  $-N(R_a)-O$ -lower alkylene, lower alkylene- $N(R_a)-O-$ ,


17 =N-O-lower alkylene, lower heteroalkylene, -O-lower heteroalkylene, -S(O)<sub>n</sub>-lower  
18 heteroalkylene, N(R<sub>a</sub>)-lower heteroalkylene, -SO<sub>2</sub>N(R<sub>a</sub>)-lower heteroalkylene, lower  
19 heteroalkylene-SO<sub>2</sub>N(R<sub>a</sub>)-, -CON(R<sub>a</sub>)-lower heteroalkylene, lower  
20 heteroalkylene-CON(R<sub>a</sub>)-, -N(R<sub>a</sub>)CON(R<sub>b</sub>)-lower heteroalkylene, lower  
21 heteroalkylene-N(R<sub>a</sub>)N(R<sub>b</sub>)-, -N(R<sub>a</sub>)SO<sub>2</sub>N(R<sub>b</sub>)-lower heteroalkylene, -N(R<sub>a</sub>)-O-lower  
22 heteroalkylene, lower heteroalkylene-N(R<sub>a</sub>)-O-, =N-O-lower alkylene, aryl and  
23 heteroaryl;

24  $L^2$  is a divalent linking group selected from the group consisting of  
25  $-\text{CH}_2\text{CH}_2-$ ,  $-(\text{C}(\text{R}_c)=\text{C}(\text{R}_d))_m-$ ,  $-\text{O}-$ ,  $-\text{S}(\text{O})_n-$ ,  $-\text{N}(\text{R}_e)-$ ,  $-\text{C}(\text{O})-$ ,  $-\text{C}(\text{O})\text{O}-$ ,  $-\text{SO}_2\text{N}(\text{R}_e)-$ ,  
26  $-\text{CON}(\text{R}_e)-$ ,  $-\text{N}(\text{R}_e)\text{CON}(\text{R}_f)-$ ,  $-\text{N}(\text{R}_e)\text{N}(\text{R}_f)-$ ,  $-\text{N}(\text{R}_e)\text{SO}_2\text{N}(\text{R}_f)-$ ,  $-\text{N}(\text{R}_e)-\text{O}-$ ,  $=\text{N}-\text{O}-$ , lower  
27 alkylene, perfluoro lower alkylene, polyfluoro lower alkylene,  $-\text{O}-$ lower alkylene,  
28  $-\text{S}(\text{O})_n-$ lower alkylene,  $\text{N}(\text{R}_e)-$ lower alkylene,  $-\text{SO}_2\text{N}(\text{R}_e)-$ lower alkylene, lower  
29 alkylene- $\text{SO}_2\text{N}(\text{R}_e)-$ ,  $-\text{CON}(\text{R}_e)-$ lower alkylene, lower alkylene- $\text{CON}(\text{R}_e)-$ ,  
30  $-\text{N}(\text{R}_e)\text{CON}(\text{R}_f)-$ lower alkylene, lower alkylene- $\text{N}(\text{R}_e)\text{N}(\text{R}_f)-$ ,  $-\text{N}(\text{R}_e)\text{SO}_2\text{N}(\text{R}_f)-$ lower  
31 alkylene,  $-\text{N}(\text{R}_e)-\text{O}-$ lower alkylene, lower alkylene- $\text{N}(\text{R}_e)-\text{O}-$ ,  $=\text{N}-\text{O}-$ lower alkylene,  
32 lower heteroalkylene,  $-\text{O}-$ lower heteroalkylene,  $-\text{S}(\text{O})_n-$ lower heteroalkylene,  $\text{N}(\text{R}_e)-$ lower  
33 heteroalkylene,  $-\text{SO}_2\text{N}(\text{R}_e)-$ lower heteroalkylene, lower heteroalkylene- $\text{SO}_2\text{N}(\text{R}_e)-$ ,  
34  $-\text{CON}(\text{R}_e)-$ lower heteroalkylene, lower heteroalkylene- $\text{CON}(\text{R}_e)-$ ,  $-\text{N}(\text{R}_e)\text{CON}(\text{R}_f)-$ lower  
35 heteroalkylene, lower heteroalkylene- $\text{N}(\text{R}_e)\text{N}(\text{R}_f)-$ ,  $-\text{N}(\text{R}_e)\text{SO}_2\text{N}(\text{R}_f)-$ lower  
36 heteroalkylene,  $-\text{N}(\text{R}_e)-\text{O}-$ lower heteroalkylene, lower heteroalkylene- $\text{N}(\text{R}_e)-\text{O}-$ ,  
37  $=\text{N}-\text{O}-$ lower alkylene, aryl and heteroaryl, wherein  $\text{R}_a$ ,  $\text{R}_b$ ,  $\text{R}_c$ ,  $\text{R}_d$ ,  $\text{R}_e$  and  $\text{R}_f$  are each  
38 members independently selected from the group consisting of H, lower alkyl, lower  
39 heteroalkyl,  $-\text{C}(\text{O})-$ lower alkyl,  $-\text{C}(\text{O})-$ lower heteroalkyl,  $-\text{S}(\text{O})_2-$ lower alkyl, and  $-\text{S}(\text{O})_2-$ lower heteroalkyl;

41 the subscript n is an integer of from 0 to 2;

42 the subscript m is an integer of from 0 to 3;

43 B is selected from the group consisting of:



wherein  $X^2$  is a substituted or unsubstituted member selected from the group consisting of a 5-6 membered cycloalkyl, 5-6 membered heterocycloalkyl containing from 1 to 3 heteroatoms, heteroaryl containing from 1 to 3 heteroatoms and aryl;

W is CH or N;

$R_g$  is selected from the group consisting of H, lower alkyl, lower alkoxy and F;


$R_h$  is selected from the group consisting of H,  $-S(O)_n$ -lower alkyl,  $-S(O)_n$ -lower heteroalkyl,  $-S(O)_n$ -aryl and  $-S(O)_n$ -heteroaryl;

$R_i$  is selected from the group consisting of H, lower alkyl, lower heteroalkyl, or a bond that links the atom bearing  $R_i$  with another atom in the  $X^2$  ring;

$R_j$  is selected from the group consisting of H, lower alkyl, F and lower alkoxy; and


Hal is a halogen atom;

wherein when  $L^1$  and  $L^2$  may be linked together *via* a single bond,  $-O-$ ,  $-S-$  or amide group to form a new 5 to 7 membered ring; with the proviso that when B is



65 and  $X^2$  is rhodanine,  $L^1$  is not  $-\text{CH}_2-\text{CH}_2-$ ,  $-\text{CH}=\text{CH}-$ ,  $-\text{C}\equiv\text{C}-$  or aryl.

1                    13.     A compound in accordance with claim 12, wherein B is selected  
2     from the group consisting of:



3  
4     wherein

5                     $R_k$  is selected from the group consisting of H, lower alkyl, lower  
6     heteroalkyl and F;

7                     $R_l$  is H or lower alkyl;


8                     $X^3$  is selected from the group consisting of O, S,  $\text{CH}_2$ ,  $\text{CH}(\text{lower alkyl})$ ,  
9      $\text{C}(\text{lower alkyl})_2$ , NH and  $\text{N}(\text{lower alkyl})$ ;

10                     $X^4$  is selected from the group consisting of O, S, NH and  $\text{N}(\text{lower alkyl})$ ,  
11     or  $X^4$  and the carbon atom to which it is attached represents an  $\text{sp}^3$ -hybridized carbon  
12     having two substituents independently selected from the group consisting of H, lower  
13     alkyl and lower heteroalkyl;

14                     $X^5$  is selected from the group consisting of O, S, NH and  $\text{N}(\text{lower alkyl})$ ,  
15     or  $X^5$  and the carbon atom to which it is attached represents an  $\text{sp}^3$ -hybridized carbon  
16     having two substituents independently selected from the group consisting of H, lower  
17     alkyl, lower alkoxy, aryloxy, lower thioalkoxy and arylthioxy; and

18                    --- represents either a single or double bond, with the proviso that when a  
19     single bond is intended, the ring atom bearing said single bond bears an additional  
20     substituent selected from the group consisting of H, lower alkyl, lower alkoxy and F.


1                    14.     A compound of claim 13, wherein B is selected from the group  
2     consisting of:



3  
4     wherein

5  $R_k$ ,  $R_m$  and  $R_n$  are each independently selected from the group consisting  
6 of H, F, lower alkyl and lower alkoxy;  
7  $X^3$  is selected from the group consisting of O, S, C(lower alkyl)<sub>2</sub>, NH and  
8 N(lower alkyl);  
9  $X^4$  is selected from the group consisting of O and S, or  $X^4$  and the carbon  
10 atom to which it is attached represents an  $sp^3$ -hybridized carbon having two substituents  
11 independently selected from the group consisting of H, lower alkyl and lower heteroalkyl;  
12  $X^5$  is selected from the group consisting of O and S, or  $X^5$  and the carbon  
13 atom to which it is attached represents an  $sp^3$ -hybridized carbon having two substituents  
14 independently selected from the group consisting of H, lower alkoxy and lower  
15 thioalkoxy.


1 **15.** A compound of claim 14, wherein B is selected from the group  
2 consisting of:



3  
4 wherein any unlabeled R groups are independently selected from the group consisting of  
5 H, lower alkyl, lower alkoxy and F.

1 **16.** A compound of claim 12 wherein B is selected from the group  
2 consisting of:






17. A compound of claim 12, wherein  $L^1$  is selected from the group consisting of  $-\text{N}(\text{R}_a)-$ ,  $-\text{N}(\text{R}_a)\text{-alkylene}$ ,  $\text{alkylene-SO}_2\text{-N}(\text{R}_a)-$ ,  $-\text{SO}_2\text{-N}(\text{R}_a)-$  and  $-\text{N}(\text{R}_a)\text{SO}_2-$ ; and  $\text{X}^1$  is selected from the group consisting of H, aryl and alkyl.


18. A compound of claim 12, wherein  $\text{Ar}^1$  is selected from the group consisting of substituted or unsubstituted biphenyl group, substituted or unsubstituted bicyclic ring, substituted or unsubstituted phenyl group and substituted or unsubstituted pyridyl.

19. A compound of claim 17, said compound having the formula:



wherein  $\text{R}_p$  is selected from the group consisting of substituted or unsubstituted alkyl, substituted or unsubstituted aryl and substituted or unsubstituted heteroaryl.

20. A compound of Claim 12, said compound having the formula (III):




wherein

the subscript  $q$  is an integer of from 0 to 4;

5  $R^1$  is hydrogen or a substituent having the formula  $-L^1-COOH$ ;  
6  $X$  is a moiety selected from  $-S-$ ,  $-O-$ , and  $-N(R_o)-$ , wherein  $R_o$  is H or  
7 lower alkyl;  
8  $R^2$  is a substituted or unsubstituted aryl( $C_1-C_8$ )alkyl, a substituted or  
9 unsubstituted aryl( $C_1-C_8$ )alkenyl, a substituted or unsubstituted aryl( $C_1-C_8$ )alkynyl, a  
10 substituted or unsubstituted alicyclic group having from 5-8 carbon atoms, or a group  
11 having the formula  $(R_{2a})_r-(L)_s-R_{2b}-$ , wherein  $R_{2a}$  and  $R_{2b}$  can be the same or different and  
12 represent a substituted or unsubstituted heterocyclic group or a substituted or  
13 unsubstituted phenyl group,  $R_{2a}$  can also represent a substituted or unsubstituted  
14 polycyclic group, and  $L$  represents a divalent linking group selected from methylene,  
15 ethylene, propylene,  $-CH=CH-$ ,  $-C\equiv C-$ ,  $-C(O)-$ ,  $-O-$ ,  $-S-$ ,  $-S(O)-$ ,  $-S(O)_2-$ , or  $-N(R_{2c})-$ ,  
16 wherein  $R_{2c}$  is selected from H or lower alkyl, and the subscripts  $r$  and  $s$  are each  
17 independently 0 or 1;  
18  $R^3$  is selected from the group consisting of H, substituted or unsubstituted  
19 ( $C_1-C_8$ )alkyl, substituted and unsubstituted aryl or substituted and unsubstituted  
20 heteroaryl;  
21  $Y$  represents O or S; and  
22  $Z$  represents O, S or  $N(R_{2d})$ , wherein  $R_{2d}$  is H or lower alkyl, or  $R_{2d}$  and  $R^1$   
23 may be joined to form an imidazole or benzimidazole group;  
24 with the proviso that when  $R^1$  is hydrogen,  $R^3$  is not substituted furan.


1 21. A compound of Claim 12, said compound having the formula (V):



2 wherein  
3  
4  $R^1$  is H,  $-OH$ ,  $-COOR_u$ ,  $-CONR_vR_w$ ,  $-SO_2NR_xR_y$  wherein  $R_u$ ,  $R_v$ ,  $R_w$ ,  $R_x$   
5 and  $R_y$  are H or lower alkyl, or  $R^1$  is a mono-heterocyclic group selected from furan,  
6 thiophene, pyridine, pyrimidine, pyridazine, 1,3-oxathiolane, tetrazole, oxadiazole,  
7 oxazole, triazole, imidazoline, imidazole, thiazole, thiadiazole, pyrrole, piperidine,  
8 morpholine, triazine and pyrazole; and  
9  $W_1$  and  $W_2$  are independently selected from H, ( $C_1-C_8$ )alkyl, ( $C_1-$   
10  $C_8$ )alkenyl, ( $C_1-C_8$ )alkynyl, halogen, nitro, hydroxy, perfluoroalkyl, difluoromethyl, ( $C_1-$

11 C<sub>8</sub>)alkoxy, phenoxy, phenyl(C<sub>1</sub>-C<sub>8</sub>)alkoxy, (C<sub>1</sub>-C<sub>8</sub>)acyl, (C<sub>1</sub>-C<sub>8</sub>)acyloxy, cyano,  
 12 carbalkoxy, thio, (C<sub>1</sub>-C<sub>8</sub>)alkylthio, (C<sub>1</sub>-C<sub>8</sub>)alkylsulfinyl, (C<sub>1</sub>-C<sub>8</sub>)alkylsulfonyl, amino, (C<sub>1</sub>-  
 13 C<sub>8</sub>)alkylamino, di(C<sub>1</sub>-C<sub>8</sub>)alkylamino, sulfonamido, carboxamido and (C<sub>1</sub>-  
 14 C<sub>8</sub>)alkanoylamino.


1                    **22.**     A compound of Claim 12, said compound having a formula  
 2 selected from the group consisting of



3  
 4 wherein


5                    R<sup>2</sup> is a substituted or unsubstituted mono- or bi-heterocyclic group, a  
 6 substituted or unsubstituted polycyclic ring, a substituted or unsubstituted alicyclic group  
 7 having 5-8 carbon atoms, a substituted or unsubstituted phenyl group, a substituted or  
 8 unsubstituted biphenyl group, a substituted or unsubstituted phenylether group, a  
 9 substituted or unsubstituted cinnameryl group, or a substituted or unsubstituted stilbenyl  
 10 group.

1                    **23.**     The compound of Claim 22, wherein said compound is selected  
 2 from the group consisting of



3  
 4

1                    **24.**     A compound having the formula (VIIa):



2

3 wherein

4  $\text{Ar}^1$  is selected from the group consisting of substituted or unsubstituted  
5 aryl and substituted or unsubstituted heteroaryl;


6 X is selected from  $-\text{S}-$ ,  $-\text{O}-$  and  $-\text{N}(\text{R}_0)-$ , wherein  $\text{R}_0$  is H or lower alkyl;

7 Y is O or S; and


8 Z is O, S or  $\text{N}(\text{R}_{2d})$ , wherein  $\text{R}_{2d}$  is H or lower alkyl, or  $\text{R}_{2d}$  and  $\text{R}^1$  may be  
9 joined to form an imidazole or benzimidazole group; and

10  $\text{R}^2$  is a substituted or unsubstituted mono- or bi-heterocyclic group, a  
11 substituted or unsubstituted polycyclic ring, a substituted or unsubstituted alicyclic group  
12 having 5-8 carbon atoms, a substituted or unsubstituted phenyl group, a substituted or  
13 unsubstituted biphenyl group, a substituted or unsubstituted phenylether group, a  
14 substituted or unsubstituted cinnamenyl group, or a substituted or unsubstituted stilbenyl  
15 group.

1 **25.** The compound of Claim 24, wherein said compound is selected  
2 from the group consisting of



1 **26.** A compound useful for the covalent modification of a viral RNA-  
2 dependent RNA polymerase (RdRp) protein, said compound having the formula:



4 wherein

5 A is an electrophilic group that reacts with a cysteine residue of said viral  
6 RNA-dependent RNA polymerase protein;

7  $\text{Ar}^1$  is a member selected from the group consisting of substituted or  
8 unsubstituted aryl and substituted or unsubstituted heteroaryl;

9  $X^1$  is a member selected from the group consisting of -H, substituted or  
10 unsubstituted alkyl, substituted or unsubstituted aryl, substituted or unsubstituted  
11 heteroaryl, -CN, -CO<sub>2</sub>H, -SO<sub>3</sub>H, -C(O)NHOH, -NH<sub>2</sub>, -OH, -NH(lower alkyl), -O(lower  
12 alkyl), -N(lower alkyl)<sub>2</sub>, and -C(O)-NH(3-tetrazolyl);

13  $L^1$  is a divalent linking group selected from the group consisting of -O-,  
14 -S(O)<sub>n</sub>-, -N(R<sub>a</sub>)-, -C(O)-, -C(O)O-, -SO<sub>2</sub>N(R<sub>a</sub>)-, -CON(R<sub>a</sub>)-, -N(R<sub>a</sub>)CON(R<sub>b</sub>)-,  
15 -N(R<sub>a</sub>)N(R<sub>b</sub>)-, -N(R<sub>a</sub>)SO<sub>2</sub>N(R<sub>b</sub>)-, -N(R<sub>a</sub>)SO<sub>2</sub>-, -N(R<sub>a</sub>)-O-, =N-O-, lower alkylene,  
16 -O-lower alkylene, -S(O)<sub>n</sub>-lower alkylene, N(R<sub>a</sub>)-lower alkylene, -SO<sub>2</sub>N(R<sub>a</sub>)-lower  
17 alkylene, lower alkylene-SO<sub>2</sub>N(R<sub>a</sub>)-, -CON(R<sub>a</sub>)-lower alkylene, lower  
18 alkylene-CON(R<sub>a</sub>)-, -N(R<sub>a</sub>)CON(R<sub>b</sub>)-lower alkylene, lower alkylene-N(R<sub>a</sub>)N(R<sub>b</sub>)-,  
19 -N(R<sub>a</sub>)SO<sub>2</sub>N(R<sub>b</sub>)-lower alkylene, -N(R<sub>a</sub>)-O-lower alkylene, lower alkylene-N(R<sub>a</sub>)-O-,  
20 =N-O-lower alkylene, lower heteroalkylene, -O-lower heteroalkylene, -S(O)<sub>n</sub>-lower  
21 heteroalkylene, N(R<sub>a</sub>)-lower heteroalkylene, -SO<sub>2</sub>N(R<sub>a</sub>)-lower heteroalkylene, lower  
22 heteroalkylene-SO<sub>2</sub>N(R<sub>a</sub>)-, -CON(R<sub>a</sub>)-lower heteroalkylene, lower  
23 heteroalkylene-CON(R<sub>a</sub>)-, -N(R<sub>a</sub>)CON(R<sub>b</sub>)-lower heteroalkylene, lower  
24 heteroalkylene-N(R<sub>a</sub>)N(R<sub>b</sub>)-, -N(R<sub>a</sub>)SO<sub>2</sub>N(R<sub>b</sub>)-lower heteroalkylene, -N(R<sub>a</sub>)-O-lower  
25 heteroalkylene, lower heteroalkylene-N(R<sub>a</sub>)-O-, =N-O-lower alkylene and heteroaryl;


26  $X^2$  is a member selected from the group consisting of substituted or  
27 unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted  
28 cycloalkyl, and substituted or unsubstituted heterocycloalkyl;

29  $L^2$  is a divalent linking group selected from the group consisting of  
30 -CH<sub>2</sub>CH<sub>2</sub>-, -(C(R<sub>c</sub>)=C(R<sub>d</sub>))<sub>m</sub>-, -O-, -S(O)<sub>n</sub>-, -N(R<sub>e</sub>)-, -C(O)-, -C(O)O-, -SO<sub>2</sub>N(R<sub>e</sub>)-,  
31 -CON(R<sub>e</sub>)-, -N(R<sub>e</sub>)CON(R<sub>f</sub>)-, -N(R<sub>e</sub>)N(R<sub>f</sub>)-, -N(R<sub>e</sub>)SO<sub>2</sub>N(R<sub>f</sub>)-, -N(R<sub>e</sub>)-O-, =N-O-, lower  
32 alkylene, perfluoro lower alkylene, polyfluoro lower alkylene, -O-lower alkylene,  
33 -S(O)<sub>n</sub>-lower alkylene, N(R<sub>e</sub>)-lower alkylene, -SO<sub>2</sub>N(R<sub>e</sub>)-lower alkylene, lower  
34 alkylene-SO<sub>2</sub>N(R<sub>e</sub>)-, -CON(R<sub>e</sub>)-lower alkylene, lower alkylene-CON(R<sub>e</sub>)-,  
35 -N(R<sub>e</sub>)CON(R<sub>f</sub>)-lower alkylene, lower alkylene-N(R<sub>e</sub>)N(R<sub>f</sub>)-, -N(R<sub>e</sub>)SO<sub>2</sub>N(R<sub>f</sub>)-lower  
36 alkylene, -N(R<sub>e</sub>)-O-lower alkylene, lower alkylene-N(R<sub>e</sub>)-O-, =N-O-lower alkylene,  
37 lower heteroalkylene, -O-lower heteroalkylene, -S(O)<sub>n</sub>-lower heteroalkylene, N(R<sub>e</sub>)-lower  
38 heteroalkylene, -SO<sub>2</sub>N(R<sub>e</sub>)-lower heteroalkylene, lower heteroalkylene-SO<sub>2</sub>N(R<sub>e</sub>)-,  
39 -CON(R<sub>e</sub>)-lower heteroalkylene, lower heteroalkylene-CON(R<sub>e</sub>)-, -N(R<sub>e</sub>)CON(R<sub>f</sub>)-lower  
40 heteroalkylene, lower heteroalkylene-N(R<sub>e</sub>)N(R<sub>f</sub>)-, -N(R<sub>e</sub>)SO<sub>2</sub>N(R<sub>f</sub>)-lower  
41 heteroalkylene, -N(R<sub>e</sub>)-O-lower heteroalkylene, lower heteroalkylene-N(R<sub>e</sub>)-O-,  
42 =N-O-lower alkylene, aryl and heteroaryl;

wherein  $R_a$ ,  $R_b$ ,  $R_c$ ,  $R_d$ ,  $R_e$  and  $R_f$  are each members independently selected from the group consisting of H, lower alkyl, lower heteroalkyl,  $-C(O)$ -lower alkyl,  $-C(O)$ -lower heteroalkyl,  $-S(O)_2$ -lower alkyl, and  $-S(O)_2$ -lower heteroalkyl; the subscript  $n$  is an integer of from 0 to 2; the subscript  $m$  is an integer of from 0 to 3; the bond between  $X^2$  and A can be a single, double or triple bond, depending on the nature of  $X^2$  and A; and wherein when  $L^1$  and  $L^2$  may be linked together *via* a single bond,  $-O-$ ,  $-S-$  or amide group to form a new 5 to 7 membered ring.

**27.** A compound in accordance with claim 26, wherein  $X^2$  is selected from the group consisting of a 5 to 7 membered cycloalkyl ring, a 5 to 7 membered heterocycloalkyl ring containing from 1 to 3 heteroatoms, an aryl group and a heteroaryl group; A is selected from the group consisting of an  $sp^2$ -hybridized carbon atom and an  $sp^3$ -hybridized carbon atom;  $L^2$  is a single bond; and  $X^2$  and A are joined *via* a single or double bond.

**28.** A compound in accordance with claim 27, wherein  $-X^2-A-$  is selected from the group consisting of:



wherein  $R_g$  is selected from the group consisting of H, lower alkyl, lower alkoxy and F;

$R_h$  is selected from the group consisting of H,  $-S(O)_n$ -lower alkyl,  $-S(O)_n$ -lower heteroalkyl,  $-S(O)_n$ -aryl and  $-S(O)_n$ -heteroaryl; W is CH or N; Hal is a halogen atom; and  $X^2$  is a substituted or unsubstituted member selected from the group consisting




31 lower heteroalkylene, -O-lower heteroalkylene, -S(O)<sub>n</sub>-lower heteroalkylene, N(R<sub>e</sub>)-lower  
 32 heteroalkylene, -SO<sub>2</sub>N(R<sub>e</sub>)-lower heteroalkylene, lower heteroalkylene-SO<sub>2</sub>N(R<sub>e</sub>)-,  
 33 -CON(R<sub>e</sub>)-lower heteroalkylene, lower heteroalkylene-CON(R<sub>e</sub>)-, -N(R<sub>e</sub>)CON(R<sub>f</sub>)-lower  
 34 heteroalkylene, lower heteroalkylene-N(R<sub>e</sub>)N(R<sub>f</sub>)-, -N(R<sub>e</sub>)SO<sub>2</sub>N(R<sub>f</sub>)-lower  
 35 heteroalkylene, -N(R<sub>e</sub>)-O-lower heteroalkylene, lower heteroalkylene-N(R<sub>e</sub>)-O-,  
 36 =N-O-lower alkylene, aryl and heteroaryl;

37 wherein R<sub>a</sub>, R<sub>b</sub>, R<sub>c</sub>, R<sub>d</sub>, R<sub>e</sub> and R<sub>f</sub> are each members independently selected  
 38 from the group consisting of H, lower alkyl, lower heteroalkyl, -C(O)-lower alkyl,  
 39 -C(O)-lower heteroalkyl, -S(O)<sub>2</sub>-lower alkyl, and -S(O)<sub>2</sub>-lower heteroalkyl;

40 the subscript n is an integer of from 0 to 2;

41 the subscript m is an integer of from 0 to 3;

42 B is selected from the group consisting of:



43 wherein

44 X<sup>2</sup> is a substituted or unsubstituted member selected from the group  
 45 consisting of a 5-6 membered cycloalkyl, 5-6 membered heterocycloalkyl containing  
 46 from 1 to 3 heteroatoms, heteroaryl containing from 1 to 3 heteroatoms and aryl;

47 W is CH or N;

48 R<sub>g</sub> is selected from the group consisting of H, lower alkyl, lower alkoxy  
 49 and F;


50 R<sub>h</sub> is selected from the group consisting of H, -S(O)<sub>n</sub>-lower alkyl, -S(O)<sub>n</sub>-  
 51 lower heteroalkyl, -S(O)<sub>n</sub>-aryl and -S(O)<sub>n</sub>-heteroaryl;

52 R<sub>i</sub> is selected from the group consisting of H, lower alkyl, lower  
 53 heteroalkyl, or a bond that links the atom bearing R<sub>i</sub> with another atom in the X<sup>2</sup> ring;



55  $R_j$  is selected from the group consisting of H, lower alkyl, F and lower  
 56 alkoxy; and  
 57 Hal is a halogen atom;  
 58 wherein when  $L^1$  and  $L^2$  may be linked together *via* a single bond, -O-, -S-  
 59 or amide group to form a new 5 to 7 membered ring.


1 **30.** A compound in accordance with claim 29, wherein B is selected  
 2 from the group consisting of:



3 wherein


5  $R_k$  is selected from the group consisting of H, lower alkyl, lower  
 6 heteroalkyl and F;  
 7  $R_1$  is H or lower alkyl;  
 8  $X^3$  is selected from the group consisting of O, S,  $CH_2$ ,  $CH$ (lower alkyl),  
 9  $C$ (lower alkyl)<sub>2</sub>, NH and  $N$ (lower alkyl);  
 10  $X^4$  is selected from the group consisting of O, S, NH and  $N$ (lower alkyl),  
 11 or  $X^4$  and the carbon atom to which it is attached represents an  $sp^3$ -hybridized carbon  
 12 having two substituents independently selected from the group consisting of H, lower  
 13 alkyl and lower heteroalkyl; and  
 14  $X^5$  is selected from the group consisting of O, S, NH and  $N$ (lower alkyl),  
 15 or  $X^5$  and the carbon atom to which it is attached represents an  $sp^3$ -hybridized carbon  
 16 having two substituents independently selected from the group consisting of H, lower  
 17 alkyl, lower alkoxy, aryloxy, lower thioalkoxy and arylthioxy; and --- represents either  
 18 a single or double bond, with the proviso that when a single bond is intended, the ring  
 19 atom bearing said single bond bears an additional substituent selected from the group  
 20 consisting of H, lower alkyl, lower alkoxy and F.

1 **31.** A compound of claim 30, wherein B is selected from the group  
 2 consisting of:




wherein  $R_k$ ,  $R_m$  and  $R_n$  are each independently selected from the group consisting of H, F, lower alkyl and lower alkoxy;  $X^3$  is selected from the group consisting of O, S, C(lower alkyl)<sub>2</sub>, NH and N(lower alkyl);  $X^4$  is selected from the group consisting of O, S, or  $X^4$  and the carbon atom to which it is attached represents an  $sp^3$ -hybridized carbon having two substituents independently selected from the group consisting of H, lower alkyl and lower heteroalkyl;  $X^5$  is selected from the group consisting of O, S, or  $X^5$  and the carbon atom to which it is attached represents an  $sp^3$ -hybridized carbon having two substituents independently selected from the group consisting of H, lower alkoxy and lower thioalkoxy.

**32.** A compound of claim 31, wherein B is selected from the group consisting of:



wherein any unlabeled R groups are independently selected from the group consisting of H, lower alkyl, lower alkoxy and F.


1 33. A compound of claim 29, wherein B is selected from the group  
2 consisting of:



1 34. A compound of claim 29 wherein  $L^1$  is selected from the group  
2 consisting of  $-N(R_a)-$ ,  $-N(R_a)$ -alkylene, alkylene- $SO_2-N(R_a)-$ ,  $-SO_2-N(R_a)-$  and  
3  $-N(R_a)SO_2-$ ; and  $X^1$  is selected from the group consisting of H, aryl and alkyl.


1 35. A compound of claim 29, wherein  $Ar^1$  is selected from the group  
2 consisting of substituted or unsubstituted biphenyl group, substituted or unsubstituted  
3 bicyclic ring, substituted or unsubstituted phenyl group and substituted or unsubstituted  
4 pyridyl.

1 36. A compound of claim 34, said compound having the formula:



2  
3 wherein  $R_p$  is a member selected from the group consisting of substituted or unsubstituted  
4 alkyl, substituted or unsubstituted aryl and substituted or unsubstituted heteroaryl.

1 37. A compound of Claim 29, said compound having the formula (III):  
2



wherein

the subscript q is an integer of from 0 to 4;

R<sup>1</sup> is hydrogen or a substituent having the formula -L<sup>1</sup>-COOH;


X is a moiety selected from -S-, -O-, and -N(R<sub>0</sub>)-, wherein R<sub>0</sub> is H or lower alkyl;

R<sup>2</sup> is a substituted or unsubstituted aryl(C<sub>1</sub>-C<sub>8</sub>)alkyl, a substituted or unsubstituted aryl(C<sub>1</sub>-C<sub>8</sub>)alkenyl, a substituted or unsubstituted aryl(C<sub>1</sub>-C<sub>8</sub>)alkynyl, a substituted or unsubstituted alicyclic group having from 5-8 carbon atoms, or a group having the formula (R<sub>2a</sub>)<sub>r</sub>-(L)<sub>s</sub>-R<sub>2b</sub>-, wherein R<sub>2a</sub> and R<sub>2b</sub> can be the same or different and represent a substituted or unsubstituted heterocyclic group or a substituted or unsubstituted phenyl group, R<sub>2a</sub> can also represent a substituted or unsubstituted polycyclic group, and L represents a divalent linking group selected from methylene, ethylene, propylene, -CH=CH-, -C≡C-, -C(O)-, -O-, -S-, -S(O)-, -S(O)<sub>2</sub>-, or -N(R<sub>2c</sub>)-, wherein R<sub>2c</sub> is selected from H or lower alkyl, and the subscripts r and s are each independently 0 or 1;

Y represents O or S; and

Z represents O, S or N(R<sub>2d</sub>), wherein R<sub>2d</sub> is H or lower alkyl, or R<sub>2d</sub> and R<sup>1</sup> may be joined to form an imidazole or benzimidazole group; with the proviso that when R<sup>1</sup> is hydrogen R<sup>2</sup> is not substituted or unsubstituted furan.

**38.** A compound of Claim 29, said compound having the formula (V):



wherein

R<sup>1</sup> is H, -OH, -COOR<sub>u</sub>, -CONR<sub>v</sub>R<sub>w</sub>, -SO<sub>2</sub>NR<sub>x</sub>R<sub>y</sub> wherein R<sub>u</sub>, R<sub>v</sub>, R<sub>w</sub>, R<sub>x</sub> and R<sub>y</sub> are H or lower alkyl, or R<sup>1</sup> is a mono-heterocyclic group selected from furan, thiophene, pyridine, pyrimidine, pyridazine, 1,3-oxathiolane, tetrazole, oxadiazole, oxazole, triazole, imidazoline, imidazole, thiazole, thiadiazole, pyrrole, piperidine, morpholine, triazine and pyrazole; and

9  $W_1$  and  $W_2$  are independently selected from H, (C<sub>1</sub>-C<sub>8</sub>)alkyl, (C<sub>1</sub>-  
10 C<sub>8</sub>)alkenyl, (C<sub>1</sub>-C<sub>8</sub>)alkynyl, halogen, nitro, hydroxy, perfluoroalkyl, difluoromethyl, (C<sub>1</sub>-  
11 C<sub>8</sub>)alkoxy, phenoxy, phenyl(C<sub>1</sub>-C<sub>8</sub>)alkoxy, (C<sub>1</sub>-C<sub>8</sub>)acyl, (C<sub>1</sub>-C<sub>8</sub>)acyloxy, cyano,  
12 carbalkoxy, thio, (C<sub>1</sub>-C<sub>8</sub>)alkylthio, (C<sub>1</sub>-C<sub>8</sub>)alkylsulfinyl, (C<sub>1</sub>-C<sub>8</sub>)alkylsulfonyl, amino, (C<sub>1</sub>-  
13 C<sub>8</sub>)alkylamino, di(C<sub>1</sub>-C<sub>8</sub>)alkylamino, sulfonamido, carboxamido and (C<sub>1</sub>-  
14 C<sub>8</sub>)alkanoylamino.


1 **39.** A compound of Claim 29, said compound having a formula  
2 selected from the group consisting of



3  
4 wherein

5  $R^2$  is a substituted or unsubstituted mono- or bi-heterocyclic group, a  
6 substituted or unsubstituted polycyclic ring, a substituted or unsubstituted alicyclic group  
7 having 5-8 carbon atoms, a substituted or unsubstituted phenyl group, a substituted or  
8 unsubstituted biphenyl group, a substituted or unsubstituted phenylether group, a  
9 substituted or unsubstituted cinnameryl group, or a substituted or unsubstituted stilbenyl  
10 group.

1 **40.** A compound having the formula:



2  
3 wherein


4 X is a member selected from the group consisting of O, S, NR<sup>11</sup> and  
5 CR<sup>11</sup>R<sup>12</sup> wherein R<sup>11</sup> and R<sup>12</sup> are each members independently selected from the group  
6 consisting of H, substituted or unsubstituted (C<sub>1</sub>-C<sub>8</sub>)alkyl, substituted or unsubstituted  
7 (C<sub>1</sub>-C<sub>8</sub>)alkoxy and substituted or unsubstituted (C<sub>1</sub>-C<sub>8</sub>)acyl;

8 Y is a member selected from the group consisting of O and S, or taken  
9 together with the carbon atom to which it is attached forms a methylene group;


10 Z<sup>1</sup> and Z<sup>2</sup> are each members independently selected from the group  
11 consisting of H and substituted or unsubstituted (C<sub>1</sub>-C<sub>8</sub>)alkoxy, or taken together form an  
12 oxo moiety;

13  $R^1$  is a member selected from the group consisting of substituted or  
 14 unsubstituted  $(C_1-C_8)$ alkyl, substituted or unsubstituted  $(C_1-C_8)$ alkylamino, substituted or  
 15 unsubstituted di $(C_1-C_8)$ alkylamino, substituted or unsubstituted  $(C_1-C_8)$ acylamino, amino,  
 16 H, substituted or unsubstituted aryl $(C_1-C_8)$ alkyl, substituted or unsubstituted  
 17 heteroaryl $(C_1-C_8)$ alkyl, substituted or unsubstituted heterocycloalkyl and  $-NHSO_2-Ar^1$ ,  
 18 wherein  $Ar^1$  is selected from the group consisting of substituted or unsubstituted aryl and  
 19 substituted or unsubstituted heteroaryl; and

20  $R^2$  and  $R^3$  are each members independently selected from the group  
 21 consisting of halo, substituted or unsubstituted  $(C_1-C_8)$ alkyl and substituted or  
 22 unsubstituted  $(C_1-C_8)$ acyl, or taken together form a group of the formula:



23  
 24 wherein  $R^4$  and  $R^5$  are each members independently selected from the  
 25 group consisting of H, substituted or unsubstituted aryl, and substituted or unsubstituted  
 26 heteroaryl, with the proviso that no more than one of  $R^4$  and  $R^5$  are H;  
 27 with the proviso that when  $Z^1$  and  $Z^2$  taken together form an oxo moiety and  $R^2$  and  $R^3$   
 28 taken together form a group of the formula:



29  
 30  $R^1$  is not substituted or unsubstituted  $(C_1-C_8)$ alkyl or H

1 **41.** A compound in accordance with claim **40**, wherein  $R^1$  is selected  
 2 from the group consisting of amino and substituted or unsubstituted  $-NHSO_2-Ar^1$ .


1 **42.** A compound in accordance with claim **40**, wherein  $Z^1$  and  $Z^2$  taken  
 2 together are oxo.

1 **43.** A compound in accordance with claim **40**, wherein Y is O or S and  
 2  $Z^1$  and  $Z^2$  taken together are oxo.

1 **44.** A compound in accordance with claim **40**, wherein X and Y are S  
 2 and  $Z^1$  and  $Z^2$  taken together are oxo.

1 **45.** A compound in accordance with claim **40**, wherein  $R^1$  is selected  
 2 from the group consisting of substituted or unsubstituted  $(C_1-C_8)$ alkylamino, substituted

3 or unsubstituted di(C<sub>1</sub>-C<sub>8</sub>)alkylamino, substituted or unsubstituted (C<sub>1</sub>-C<sub>8</sub>)acylamino,  
4 amino, and -NHSO<sub>2</sub>-Ar<sup>1</sup>, wherein Ar<sup>1</sup> is selected from the group consisting of substituted  
5 or unsubstituted aryl and substituted or unsubstituted heteroaryl; X and Y are each  
6 independently selected from the group consisting of O and S; Z<sup>1</sup> and Z<sup>2</sup> taken together are  
7 oxo; and R<sup>2</sup> and R<sup>3</sup> taken together are a group having the formula:



8  
9 wherein R<sup>4</sup> and R<sup>5</sup> are each members independently selected from the  
10 group consisting of H, substituted or unsubstituted aryl, and substituted or unsubstituted  
11 heteroaryl, with the proviso that only one of R<sup>4</sup> and R<sup>5</sup> is H.

Sub  
B1  
1 46. A pharmaceutical composition, comprising a pharmaceutically  
2 acceptable carrier and a therapeutically or prophylactically effective amount of a  
3 compound of any one of claims 1, 9, 12, 24, 26, 29 and 40.

1 47. A method for the treatment or prevention of a viral infection,  
2 comprising  
3 administering to a subject suffering from or at risk for said viral infection  
4 an effective amount of a compound of any one of claims 1, 9, 12, 24, 26, 29 and 40.

1 48. The method of Claim 47, wherein said viral infection is hepatitis C  
2 virus infection.

1 49. The method of Claim 47, wherein said compound is administered  
2 in combination with a therapeutically effective amount of an antiviral agent.

1 50. The method of Claim 49, wherein said antiviral agent is an  
2 interferon.

1 51. A method for treating or preventing a viral infection, comprising  
2 administering to a subject in need thereof a therapeutically effective  
3 amount of a compound that binds to a cysteine residue in the RNA-dependent RNA  
4 polymerase (RdRp) protein of a virus forming a covalent bond.

1 52. The method of Claim 51, wherein said RdRp protein is NS5B.

1                   **53.**     The method of Claim **51**, wherein said viral infection is hepatitis C  
2 virus infection.

1                   **54.**     The method of Claim **51**, wherein said compound comprises an  
2 electrophilic group that reacts with a cysteine residue of said RdRp protein.

1                   **55.**     The method of Claim **54**, wherein said electrophilic group is  
2 selected from the group consisting of an activated double or triple bond, an electrophilic  
3 center, a carboxylic acid or carboxylic acid derivative, a sulfur-containing group and an  
4 activated or unactivated carbonyl group.

0938270 040501  
"05040" 0232960